

Flux Quantization Due to Monopole and Dipole Currents

Wei Chen,¹ Peter Horsch,¹ and Dirk Manske^{1,2}

¹Max-Planck-Institut für Festkörperforschung, Heisenbergstrasse 1, D-70569 Stuttgart, Germany

²Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan

(Dated: February 5, 2013)

We demonstrate the general principle for flux quantization in field-induced quantum interference effects. For particles carrying a monopole moment, the interference causes monopole current to oscillate periodically with flux defined as inner product of field and area, whereas for particles carrying a fixed dipole moment, the dipole current oscillates periodically with flux vector defined as cross product of field and trajectory. Our analysis unifies the oscillation of monopole or dipole currents in various devices, such as SQUID and spin-FET, into the same physical picture. As consequence of these general principles and the non-gauge field nature of the coupling between dipole moments and external field, it follows that the quantization of the electric flux vector can also be seen in open trajectory devices that transport dipole currents, such as spin Josephson effect.

PACS numbers: 03.65.Vf, 03.75.Lm, 85.35.Ds, 85.75.Hh, 14.80.Hv

Introduction.- Quantum interference effects due to external electric or magnetic fields have long been of great interest, since they demonstrate the feasibility of controlling the quantum state of particles by external fields and preserving quantum coherence. Such interference effects stem from the coupling of monopole or dipole moment of the particles to the vector field or the electric and magnetic field via Aharonov-Bohm (AB)[1], dual Aharonov-Bohm (DAB)[2], Aharonov-Casher (AC)[3], and He-McKellar-Wilkens effects (HMW)[4, 5]. Due to these effects, the wave function of the particle acquires a Berry phase after traveling along a certain trajectory, and allows to utilize the Berry phase to design various interferometers.

Since the Berry phase is a periodic argument, a natural consequence of field-induced interference effects is the flux quantization. A well-known example is the quantization of magnetic flux in a superconducting quantum interference device (SQUID), in which the Berry phase due to AB effect combined with single-valuedness of the wave function yields a Josephson current oscillating periodically with magnetic flux, from which the flux quantization is interpreted. The magnetic flux is defined as the inner product of magnetic field and the area enclosed by the trajectory; and the smallness of the flux quantum $h/2e$ is the reason behind the high precision magnetometer made by SQUID.

General principle of flux quantization.- In this Letter, we show that flux quantization can also be introduced in DAB, AC, and HMW effects in the same sense as flux quantization in a SQUID, i.e., from the oscillation of monopole or dipole currents with flux. We follow the philosophy that the flux should be defined as the quantity that controls the interference effect but only depends on the field and the trajectory, whereas the monopole or dipole moments of the particles determine the flux quantum. The oscillation of monopole or dipole currents due to quantum interference is best demonstrated

within the framework of persistent current in a mesoscopic ring[6, 7], in combination with the generic set up proposed in Ref. 2. As duplicated in Fig. 1(a), quantum particles carrying electric charge q /magnetic monopole q_m / magnetic dipole $\boldsymbol{\mu}$ /electric dipole $\mathbf{d}$ are confined in a 1D ring of length L , with an infinitely long wire carrying uniformly distributed magnetic dipole $\boldsymbol{\mu}$ /electric dipole $\mathbf{d}$ /electric charge q /magnetic monopole q_m pierced through the center, which corresponds to the set up for AB/DAB/AC/HMW effects, respectively. First we review the mechanism of persistent charge current due to AB effect in this set up, then we make analogy to the other three effects.

In the set up for AB effect, the momentum $\mathbf{p}$ of a particle with electric charge q is replaced by (SI units are adopted throughout the article, and boldface symbols are vectors)

$$AB: \quad \mathbf{p} - q\mathbf{A} \rightarrow \varphi_{AB} = \frac{q}{\hbar} \oint \mathbf{A} \cdot d\mathbf{l} = 2\pi \frac{\Phi_B}{\Phi_B^0}, \quad (1)$$

that is, from the form of momentum, the Berry phase φ_{AB} is already determined. The discrete eigenenergies for the particles with mass m are

$$E_n = \frac{2\pi^2 \hbar^2}{mL^2} (n - \Phi_B/\Phi_B^0)^2. \quad (2)$$

The many particle energy spectrum is periodic in $\Phi_B = \oint \mathbf{A} \cdot d\mathbf{l} = \int \mathbf{B} \cdot d\mathbf{a}$ with periodicity $\Phi_B^0 = h/q$, and so is the charge current calculated by $I = \sum_n (q/\hbar L) f(E_n) \partial E_n / \partial k$, where $f(E_n)$ is the Fermi or Bose distribution function depending on the statistics of the particles[8]. We emphasize that although the AB effect is due to coupling of q to the vector field $\mathbf{A}$, the magnetic flux is defined only through the identity $\oint \mathbf{A} \cdot d\mathbf{l} = \int \mathbf{B} \cdot d\mathbf{a}$.

Now consider the set up for the DAB effect. Following Ref. 2, the electric field due to $\mathbf{d}$ distributed on the wire

is written in terms of a vector field $\mathbf{E} = \nabla \times \mathbf{A}_E$, and the momentum of the particle that carries q_m is replaced by

$$\begin{aligned} DAB : \quad & \mathbf{p} + \frac{q_m}{c^2} \mathbf{A}_E \\ \rightarrow \quad & \varphi_{DAB} = -\frac{q_m}{\hbar c^2} \oint \mathbf{A}_E \cdot d\mathbf{l} = 2\pi \frac{\Phi_E}{\tilde{\Phi}_E^0} \end{aligned} \quad (3)$$

where $\Phi_E = -\oint \mathbf{A}_E \cdot d\mathbf{l} = -\int \mathbf{E} \cdot d\mathbf{a}$, and $\tilde{\Phi}_E^0 = \hbar c^2/q_m$. The eigenenergies have the same form as Eq. (2), with the replacement $\Phi_B/\Phi_B^0 \rightarrow \Phi_E/\tilde{\Phi}_E^0$. Following the same argument as in the AB effect, both the eigenenergies and monopole current are periodic in Φ_E with periodicity $\tilde{\Phi}_E^0$, hence the quantization of Φ_E is interpreted.

It is intriguing to ask if our analysis can be applied to the recently discovered magnetic monopoles in spin ice [9–11] or quantum spin ice [12, 13]. For instance, if piercing an $\mathbf{E}$ field through a ring made of quantum spin ice can generate a monopole current that oscillates with Φ_E . However, it is clear that in our analysis, we assume an ideal monopole whose presence modifies Maxwell's equation, and it experiences a Lorentz force in the presence of an $\mathbf{E}$ field. Since the monopoles in quantum spin ice do not satisfy such criteria, the DAB effect does not occur in these materials. Therefore our analysis for such ideal monopoles is rather for the sake of completing the duality between magnetism and electricity, whereas its realization depends on the existence of ideal monopoles.

Similarly, for the AC and HMW effects; if charge neutral particles carry a fixed dipole moment

$$\begin{aligned} AC : \quad & \mathbf{p} - \frac{1}{c^2} \mathbf{E} \times \boldsymbol{\mu} \rightarrow \varphi_{AC} = \frac{1}{\hbar c^2} \int (\mathbf{E} \times \boldsymbol{\mu}) \cdot d\mathbf{l} \\ & = \frac{|\boldsymbol{\mu}|}{\hbar c^2} \hat{\boldsymbol{\mu}} \cdot \left(\int d\mathbf{l} \times \mathbf{E} \right) = 2\pi \frac{\hat{\boldsymbol{\mu}} \cdot \boldsymbol{\Phi}_E}{\tilde{\Phi}_E^0}, \\ HMW : \quad & \mathbf{p} - \mathbf{d} \times \mathbf{B} \rightarrow \varphi_{HMW} = \frac{1}{\hbar} \int (\mathbf{d} \times \mathbf{B}) \cdot d\mathbf{l} \\ & = \frac{|\mathbf{d}|}{\hbar} \hat{\mathbf{d}} \cdot \left(\int \mathbf{B} \times d\mathbf{l} \right) = 2\pi \frac{\hat{\mathbf{d}} \cdot \boldsymbol{\Phi}_B}{\tilde{\Phi}_B^0}. \end{aligned} \quad (4)$$

The Berry phase again is deduced directly from the form of the momentum. By factoring the Berry phase into the part that depends only on the field and trajectory and the part that depends on the fixed dipole moment, the flux is unavoidably a vector defined by the cross product of field and trajectory. Below we call $\boldsymbol{\Phi}_E$ the electric flux vector, and $\boldsymbol{\Phi}_B$ the magnetic flux vector. The eigenenergies follow Eq. (2), with the replacement $\Phi_B/\Phi_B^0 \rightarrow \hat{\boldsymbol{\mu}} \cdot \boldsymbol{\Phi}_E/\tilde{\Phi}_E^0$ and $\Phi_B/\Phi_B^0 \rightarrow \hat{\mathbf{d}} \cdot \boldsymbol{\Phi}_B/\tilde{\Phi}_B^0$ for AC and HMW effect, respectively. Both the eigenenergies and dipole currents are periodic in $\boldsymbol{\Phi}_E$ and $\boldsymbol{\Phi}_B$, from which their quantization follows. The actual quantized values of $\boldsymbol{\Phi}_E$ and $\boldsymbol{\Phi}_B$ depend not only on their flux quanta $\tilde{\Phi}_E^0 = \hbar c^2/|\boldsymbol{\mu}|$ and $\tilde{\Phi}_B^0 = \hbar/|\mathbf{d}|$, but also on the direction of the dipole moments. The oscillation of dipole currents with field has been discussed by Balatsky and Altshuler, who considered persistent spin currents in a ^3He ring[15], and the

	electric	magnetic
monopole	$\Phi_B = \int \mathbf{B} \cdot d\mathbf{a},$ $\Phi_B^0 = \hbar/q$	$\Phi_E = -\int \mathbf{E} \cdot d\mathbf{a},$ $\Phi_E^0 = \hbar c^2/q_m$
dipole	$\boldsymbol{\Phi}_B = \int \mathbf{B} \times d\mathbf{l},$ $\tilde{\Phi}_B^0 = \hbar/ \mathbf{d} $	$\boldsymbol{\Phi}_E = \int d\mathbf{l} \times \mathbf{E},$ $\tilde{\Phi}_E^0 = \hbar c^2/ \boldsymbol{\mu} $

TABLE I: Proper definition of flux and flux quantum in the field-induced interference effects, classified according to the fixed monopole or dipole moment of the particles.

electric flux quantum $\tilde{\Phi}_E^0$ has been introduced by Bogachev and Landman[14]. While these considerations refer to flux quantization in closed circuits, our point is that this oscillation obeys a general principle of flux quantization where the proper definition of flux is a vector for fixed dipole moments. Next, we address how quantization of flux vectors can be realized in concrete devices.

Realizations of quantization of electric flux vector. - For particles that carry electron magnetic moment μ_B and experience AC effect, the quantum for electric flux vector $\boldsymbol{\Phi}_E$ is huge $\tilde{\Phi}_E^0 = \hbar c^2/\mu_B = 6.43 \times 10^6 \text{V}$. This means for a reasonable scale of $\mathbf{E}$ field, φ_{AC} is only of the order of mrad[16]. In comparison with the smallness of magnetic flux quantum $\Phi_B^0 = \hbar/2e = 2.07 \times 10^{-15} \text{Wb}$, one may wonder if such a huge flux quantum and the resulting small phase shift can be of any use. Below we demonstrate that in devices that contain Rashba spin-orbit coupling (SOC), which yields a phase shift also proportional to $\mathbf{E} \times \boldsymbol{\mu}$ but with a much larger prefactor, the flux quantum may be experimentally observable.

A prototype interference device that utilizes Rashba SOC is the spin field effect transistor (spin-FET)[17]. In an ideal 1D spin-FET with length L along x -direction, the spin degeneracy ($\sigma = \pm 1$) is lifted in the 2DEG region where an $\mathbf{E}$ field is applied along $\hat{\mathbf{y}}$ direction, described by $E_\sigma = \hbar^2 k_{x\sigma}^2/2m^* - \sigma \alpha k_{x\sigma}$. In presence of Rashba coupling the motion along x -direction is then described by the Hamiltonian $H_\sigma = |\mathbf{p}_\sigma - \sigma(\mathbf{g} + \lambda \mathbf{E}) \times \boldsymbol{\mu}|^2/2m - |(\mathbf{g} + \lambda \mathbf{E}) \times \boldsymbol{\mu}|^2/2m$. The Rashba coupling corresponds to $\alpha = \hbar |(\mathbf{g} + \lambda \mathbf{E}) \times \boldsymbol{\mu}|/m$, where $\mathbf{g}$ represents the intrinsic SOC of the 2DEG, and λ characterizes the field induced SOC. It is still debated whether the field dependence of α comes from the expectation value of the electric field at the interface[18], or the asymmetry of the wave function in the quantum well[19, 20], or other aspects of the wave function[21]. For either mechanism to be true, our point is that the Rashba parameter can be *empirically* written in the form $\alpha = \hbar |(\mathbf{g} + \lambda \mathbf{E}) \times \boldsymbol{\mu}|/m$ where $\mathbf{E}$ is the external field depending only on the gate voltage and sample thickness. By preparing the spins in the source and drain in an eigenstate of σ_x , the tunneling

probability $P_0 \propto 1 + \cos \Delta\theta$ oscillates with

$$\begin{aligned}\Delta\theta &= (k_{x+} - k_{x-})L = \frac{2m\eta L}{\hbar^2} = \int (\mathbf{k}_+ - \mathbf{k}_-) \cdot d\mathbf{l} \\ &= \varphi_0 + \frac{2\lambda|\boldsymbol{\mu}|}{\hbar} \hat{\boldsymbol{\mu}} \cdot \int d\mathbf{l} \times \mathbf{E} = \varphi_0 + 2\pi \frac{\hat{\boldsymbol{\mu}} \cdot \boldsymbol{\Phi}_E}{\tilde{\Phi}_E^0}. \quad (5)\end{aligned}$$

Therefore from the periodicity of tunneling probability, or equivalently the current-voltage characteristics, the quantization of $\boldsymbol{\Phi}_E$ is realized, with flux quantum $\tilde{\Phi}_E^0 = h/2\lambda|\boldsymbol{\mu}|$. Physically, the quantization of $\boldsymbol{\Phi}_E$ shows up because spin-FET utilizes the Berry phase of spin up and down (which are fixed magnetic moments) to control the tunneling probability, so it falls into the category of field-induced quantum interference effect. Applying a gate voltage $\sim 1\text{V}$ on an inverted $\text{In}_{0.53}\text{Ga}_{0.47}\text{As}/\text{In}_{0.52}\text{Al}_{0.48}\text{As}$ heterostructure of thickness $\sim 100\text{nm}$ and channel length $\sim \mu\text{m}$ one can obtain $\Delta\theta = 2\pi[18]$, which corresponds to a flux quantum $\tilde{\Phi}_E^0 \sim 10\text{V}$. Thus the flux quantum due to Rashba SOC is generally much smaller and more accessible compared to that in AC effect.

Flux quantization in open trajectory devices.— Although we discuss the flux quantization due to monopole currents and due to dipole currents on equal footing, a crucial difference between them should be emphasized. The quantization due to monopole current originates from the coupling of monopole moment and the vector field $\mathbf{A}$ and $\mathbf{A}_E$ described in Eqs. (1) and (3), which can always be gauged away unless the particle is moved in a closed trajectory, such that the gauge invariant fluxes $\oint \mathbf{A} \cdot d\mathbf{l} = \int \mathbf{B} \cdot d\mathbf{a}$ and $-\oint \mathbf{A}_E \cdot d\mathbf{l} = -\int \mathbf{E} \cdot d\mathbf{a}$ can be measured. On the other hand, the quantization due to dipole current stems from the coupling of dipole moment and the external field directly, which cannot be gauged away even if the particle moves in an open trajectory, as demonstrated in Fig. 1(b). In other words, AC and HMW effects can cause oscillation of dipole current in open trajectory devices.

An example of such open trajectory devices is the Josephson junction. This motivates us to study spin Josephson effect in the presence of an electric field. In particular, we revisit the spin Josephson effect due to coherent tunneling of spinful particle-hole pair $\langle c_{\uparrow}^{\dagger} c_{\downarrow} \rangle$ in a ferromagnetic metal/insulator/ferromagnetic metal (FMM/I/FMM) junction,[22] instead of tunneling of spinful Cooper pairs[23] because the charge neutral $\langle c_{\uparrow}^{\dagger} c_{\downarrow} \rangle$ couples to external electric field via SOC or AC effect, and there is no Josephson charge current in this problem. We follow the analysis in Ref. 22 for FMM/I/FMM junction but with a different mean field treatment of Hubbard interaction in bulk FMM

$$Un_{\uparrow}(\mathbf{r})n_{\downarrow}(\mathbf{r}) \rightarrow -c_{\uparrow}^{\dagger}(\mathbf{r})c_{\downarrow}(\mathbf{r})\Delta(\mathbf{r}) - c_{\downarrow}^{\dagger}(\mathbf{r})c_{\uparrow}(\mathbf{r})\Delta(\mathbf{r})^{\dagger} \quad (6)$$

where $\Delta^{\dagger} = U\langle S^+ \rangle = U\langle S^x + iS^y \rangle = U\langle c_{\uparrow}^{\dagger} c_{\downarrow} \rangle$. Thus our magnetization lies in the $S^x S^y$ -plane with an angle

θ that also determines the phase of $\Delta = |\Delta|e^{i\theta}$, hence the dynamics of magnetization is directly related to the uncertainty relation $[\theta, S^z] = i$.

We can directly utilize the formalism in Ref. 22 to calculate the spin current by setting $m = (2U/3)\langle S^z \rangle = 0$ therein. Consider that the magnetizations on the two sides have the same magnitude $|\Delta_L| = |\Delta_R| = |\Delta|$ but a difference in direction $\theta = \theta_L - \theta_R$. Then the spin supercurrent due to coherent tunneling of $\langle c_{\uparrow}^{\dagger} c_{\downarrow} \rangle$ is

$$\begin{aligned}J_s &= \frac{1}{2} \langle \dot{N}_{L\uparrow} - \dot{N}_{L\downarrow} \rangle = \frac{|T|^2}{2} S(0, |\Delta|) \sin \theta \\ &= J_s^0 \sin \theta = N \langle \dot{S}_L^z \rangle = -N \langle \dot{S}_R^z \rangle, \quad (7)\end{aligned}$$

where $|T|^2$ represents the tunneling amplitude, and N is total number of sites on either side of the junction. The function $S(a, b)$ satisfies $S(a, 0) = 0$ and $S(0, b) \neq 0$.




FIG. 1: (color online) (a) Generic set up that manifests oscillation of monopole or dipole currents due to AB, DAB, AC, and HMW effects. (b) Schematics of the open trajectory devices in which AC and HMW effects can take place. (c) Proposed 2DFMM/2DEG/2DFMM junction as a realization of the open trajectory device sketched in (b). Magnetization $\langle \mathbf{S}_R \rangle$ and $\langle \mathbf{S}_L \rangle$ (blue arrows) yield the Josephson spin current J_s (polarization $\boldsymbol{\mu} \parallel \langle \mathbf{S}_R \times \mathbf{S}_L \rangle$) described by Eq. (10). The coplanar magnetization profile at the 2DEG interface under the influence of field-induced SOC (red arrows) deviates from the case without SOC (green arrows) by the phase shift φ_{AC} . Here φ_{AC} of the φ -junction can be controlled by the gate voltage and may be measured by the optical Kerr effect.

Applying a gate voltage on the thin insulating interface has two effects. Firstly, it changes the potential barrier at the interface, thus changing the tunneling amplitude $|T|^2$. Therefore one should replace $J_s^0 \rightarrow J_s^0(\mathbf{E})$. Secondly, the propagation of spinful $\langle c_{\uparrow}^{\dagger} c_{\downarrow} \rangle$ picks up a phase that depends linearly on the electric field due to AC effect or Rashba SOC. From previous calculation

for spin-FET, we anticipate that strong Rashba SOC is also necessary to experimentally observe flux quantum in spin Josephson effect. This leads us to consider the 2DFMM/2DEG/2DFMM junction shown in Fig.1(c), described by the following Ginzburg-Landau (GL) free energy that contains an AC phase

$$f = f_{n0} + \alpha|\psi|^2 + \frac{\beta}{2}|\psi|^4 - \frac{\hbar^2}{2m} |(\partial_x - ik_{0x})\psi|^2 \quad (8)$$

where α and β are GL parameters, m represents the effective mass, $\mathbf{k}_0 = (\mathbf{g} + \lambda\mathbf{E}) \times \boldsymbol{\mu}/\hbar$ and k_{0x} is its component along the junction. Eq. (8) is consistent with the Hamiltonian of a spin current in an electric field [24], with a fixed dipole moment $\boldsymbol{\mu} \parallel \langle \mathbf{S}_R \times \mathbf{S}_L \rangle$. The $\mathbf{g}$ and λ again represent intrinsic and field dependence of Rashba SOC at the interface. The dimensionless quantity $g(x) = \psi(x)/\psi_\infty = \psi_x/(-\alpha/\beta)$ in the interface $0 < x < L$ where the $\mathbf{E}$ field is applied is

$$g(x) = (1 - \frac{x}{L})e^{ik_{0x}x} + \frac{x}{L}e^{-ik_{0x}(L-x)+i\theta} \quad (9)$$

such that it satisfies the boundary condition $g(0) = 1$ and $g(L) = e^{i\theta}$, and the Laplace equation $(\partial_x - ik_{0x})^2 g = 0$. [25] The GL free energy integrated over the interface is $\Delta F = l_c \hbar^2 \psi_\infty^2 / Lm [1 - \cos(\theta - k_{0x}L)]$, where l_c is the cross section length. The current from Eq. (8) yields

$$J_s = J_s^0(\mathbf{E}) \sin(\theta - \varphi_0 - \varphi_{AC}) \quad (10)$$

where $\varphi_0 = \boldsymbol{\mu} \cdot \int d\mathbf{l} \times \mathbf{g}/\hbar$, and $\varphi_{AC} = (\lambda|\boldsymbol{\mu}|/\hbar) \hat{\boldsymbol{\mu}} \cdot \int d\mathbf{l} \times \mathbf{E} = 2\pi \hat{\boldsymbol{\mu}} \cdot \boldsymbol{\Phi}_E / \tilde{\Phi}_E^0$. The periodicity of Eq. (10) again implies $\boldsymbol{\Phi}_E$ is quantized by $\tilde{\Phi}_E^0 = h/\lambda|\boldsymbol{\mu}|$.

In the absence of Rashba SOC, $k_{0x} = 0$, Eq.(9) implies the magnetization profile in the 2DEG interface (green arrows in Fig.1(c)) gradually rotates from $\langle \mathbf{S}_L \rangle$ to $\langle \mathbf{S}_R \rangle$, due to its proximity to the two FMMs. Gradient of this coplanar magnetization is the origin of the Josephson spin current that has a fixed polarization $\boldsymbol{\mu} \parallel \langle \mathbf{S}_R \times \mathbf{S}_L \rangle$. The Rashba SOC gives additional rotation to the magnetization profile (red arrows in Fig.1(c)). Hence the gate voltage changes the magnetization profile in the interface, which should be visible as certain fringe pattern by optical Kerr effect. Details of this 2D junction will be presented in a forthcoming work.

Regarding quantization of magnetic flux vector.- An intense investigation has been dedicated to the realization of HMW effect. Since there is no sizeable $\mathbf{d}$ for known point particles, existing proposals mainly focus on field induced electric dipole moments [27, 28]. Recently, interference effect due to electrically polarized ^7Li ions has been observed in an atom interferometer [29], with an interferometer signal $I = I_0 (1 + \mathcal{V} \cos(\varphi_p + \varphi_d))$ that depends on fringe visibility $\mathcal{V}$, perturbation phase φ_p , and diffraction phase φ_d . Following previous arguments for spin-FET and spin Josephson effect, it is tempting to directly associate the periodicity of I to quantization of

$\boldsymbol{\Phi}_B$. However, the complication of such electrically polarized atom is that, by travelling through a region with both $\mathbf{E}$ and $\mathbf{B}$ field, the atom picks up a phase not only due to HMW effect, but also due to AC, Zeeman, and Stark effects. Further treatment is necessary to extract the HMW phase. Therefore one cannot directly attribute the periodicity to flux quantization. To actually interpret flux quantization from oscillation of electric dipole current or certain interference pattern, a design that only HMW phase contributes to the interference is necessary. Such design remains unclear to us at this moment.

Conclusions.- We formulate the general principle for flux quantization in field-induced quantum interference effects. By factoring the Berry phase into the part that depends only on the field and trajectory and the part that depends on the monopole or dipole moment, the flux is a scalar for particles carrying a monopole moment, but a vector for particles carrying a fixed dipole moment. This principle unifies all field-induced quantum interference devices that transport fixed monopole or dipole moments, including already known examples such as SQUID, spin-FET, and persistent charge or spin current in a mesoscopic ring. Due to the non-gauge field character of the coupling between dipole moments and the external field, we predict that such quantization can also be seen in open trajectory devices that transport dipole currents, such as spin Josephson effect. In particular we show that because of the reduced flux quantum in systems with Rashba SOC, electric flux quantization may become accessible to experiment.

We thank M. Sigrist, P. M. R. Brydon, O. P. Sushkov, H. Nakamura, W. Metzner, and F. S. Nogueira for stimulating discussions.

-
- [1] Y. Aharonov and D. Bohm, Phys. Rev. **115**, 485 (1959).
 - [2] J. P. Dowling, C. P. Williams, and J. D. Franson Phys. Rev. Lett. **83**, 2486 (1999).
 - [3] Y. Aharonov and A. Casher, Phys. Rev. Lett. **53**, 319 (1984).
 - [4] X.-G. He and B. H. J. McKellar, Phys. Rev. A **47**, 3424 (1993).
 - [5] M. Wilkens, Phys. Rev. Lett. **72**, 5 (1994).
 - [6] N. Byers and C. N. Yang, Phys. Rev. Lett. **7**, 46 (1961).
 - [7] M. Büttiker, Y. Imry, and R. Landauer, Phys. Lett. **96**, 365 (1983).
 - [8] H.-F. Cheung, Y. Gefen, E. K. Riedel, and W.-H. Shih, Phys. Rev. B **37**, 6050 (1988).
 - [9] C. Castelnovo, R. Moessner and S. L. Sondhi, Nature **451**, 42 (2008).
 - [10] D. J. P. Morris *et al.*, Science **326**, 411 (2009).
 - [11] S. Bramwell *et al.*, Nature **461**, 956 (2009).
 - [12] M. Hermele, M. P. A. Fisher, and L. Balents, Phys. Rev. B **69**, 064404 (2004).
 - [13] L. Savary and L. Balents, Phys. Rev. Lett. **108**, 037202 (2012).
 - [14] E. N. Bogachev and U. Landman, Phys. Rev. B **50**, 2678

- (1994).
- [15] A. V. Balatsky and B. L. Altshuler, Phys. Rev. Lett. **70**, 1678 (1993).
 - [16] A. Cimmino *et al.*, Phys. Rev. Lett. **63**, 380 (1989).
 - [17] S. Datta and B. Das, Appl. Phys. Lett. **56**, 665 (1990).
 - [18] J. Nitta, T. Akazaki, H. Takayanagi, and T. Enoki, Phys. Rev. Lett. **78**, 1335 (1997).
 - [19] G. Engels, J. Lange, Th. Schäpers, and H. Lüth, Phys. Rev. B **55**, 1958 (1997).
 - [20] W. Zawadzki and P. Pfeffer, Semicond. Sci. Technol. **19**, R1-R17 (2004).
 - [21] D. Grundler, Phys. Rev. Lett. **84**, 6074 (2000).
 - [22] F. S. Nogueira and K.-H. Bennemann, Europhys. Lett. **67**, 620 (2004).
 - [23] Y. Asano, Phys. Rev. B **74**, 220501(R) (2006).
 - [24] F. Meier and D. Loss, Phys. Rev. Lett. **90**, 167204 (2003).
 - [25] M. Tinkham, *Introduction to Superconductivity*, McGraw-Hill (1996), p. 199.
 - [26] V. Ambegaokar and A. Baratoff, Phys. Rev. Lett. **10**, 486 (1963); *ibid.* **11**, 104 (1963).
 - [27] H. Wei, R. Han, and X. Wei, Phys. Rev. Lett. **75**, 2071 (1995).
 - [28] Y. Sato and R. Packard, J. Phys. Conf. Ser. **150**, 032093 (2009).
 - [29] S. Lepoutre, A. Gauguier, G. Trénec, M. Büchner, and J. Vigué, Phys. Rev. Lett. **109**, 120404 (2012).